



1 in 75 That's how many women will be diagnosed with ovarian cancer during their lifetime.

Although ovarian cancer is typically associated with women after menopause, those with mucinous ovarian cancer (MOC) are often diagnosed under the age of 40.



YOU'RE NOT OVARY-ACTING!



If you feel as if your doctor isn't listening, be persistent or find a new doctor. It's not uncommon for women to be misdiagnosed, especially in the early stages.

There is no screening test for ovarian cancer.

DID YOU KNOW?

Your yearly visit with your gynecologist is important, but doesn't help identify ovarian cancer.

Be proactive. Ask your gynecologist about personal risk factors. Learn the early symptoms.

Be Aware of Common Symptoms.

- Bloating
- Back pain
- Indigestion
- Fatigue
- Change in appetite
- More frequent urination
- Abdominal pain/pressure

If these symptoms persist two weeks or longer, call your doctor.



Join us in the fight by
making an online donation

SCAN ME



Despite being the leading cause of gynecologic cancer deaths among women and the fifth leading cause of cancer deaths overall, research for ovarian cancer is woefully underfunded.

At the Mucinous Ovarian Cancer Coalition we are dedicated to improving the survival rate and quality of life for each of these women and the people who love them. As a 501(c)(3) organization, we strive to replace fear and pain with hope, comfort, and joy.

STAY CONNECTED:



hope4moc.org



[@MucinousOvarianCancerCoalition](https://www.facebook.com/MucinousOvarianCancerCoalition)



Hope4MOC@gmail.com

#HerFightIsOurFight • #Hope4MOC • #MucinousOvarianCancer